Tips to Increase 4th DTaP Rates

Parents need to hear from you about the importance of the 4th DTaP dose, which is often missed. When you speak, parents listen. Pertussis rates in Washington State are significantly higher than the national average. Many children in your practice may be at risk if they don't receive the 4th DTaP on time. Make sure your patients get the 4th DTaP dose at 15-18 months.

Use these tips to ensure children receive all five DTaP doses:

- Recommend vaccination even when a child has a mild illness, such as a cold, diarrhea or an ear infection.
- Remember: it is safe to give multiple vaccines at every visit.
- Review a child's immunization history at every visit, including acute visits.
- Record immunizations given in the CHILD Profile Immunization Registry.
- Use reminder/recall cards to ensure parents bring children in for each DTaP dose. Use the CHILD Profile Immunization Registry or order free Reminder/Recall Cards (information on back).
- Give the 3rd DTaP dose on time. Research shows that children who receive the 3rd dose on time are more likely to receive the 4th dose on time.
- Pre-schedule 4th DTaP appointments when children receive their 3rd DTaP, if possible.
- Find out your practice's 4th DTaP coverage rates and track them over time. Ask your local health department how they can help you do this.

Be part of the solution. Get involved!

Help Washington fight pertussis and increase 4th DTaP immunization rates:

- Contact your local health department to:
 - set up an assessment to review your practice's immunization coverage
 - get materials to educate your staff
 - find additional immunization information
- **Visit** the Washington State Department of Health website to:
 - order a free 4th DTaP poster for your office
 - order Reminder/Recall Cards www.doh.wag.gov/cfh/immunize/formpubs.htm
- **Join** the Immunization Action Coalition of Washington at www.withinreachwa.org/iacw

Follow the Advisory Committee on Immunization Practices recommendations for DTaP vaccination:

Ist DTaP, 2 months 2nd DTaP, 4 months 3rd DTaP, 6 months

4th DTaP, 15-18* months

5th DTaP, 4-6 years

When you speak, parents listen.

Make sure your patients don't miss any DTaP doses.





PUBLIC HEALTH
ALWAYS WORKING FOR A SAFER AND
HEALTHIER WASHINGTON

This publication is available in other formats for people with disabilities. Call 1-800-525-0127 or TTY relay 711.

^{*}The 4th DTaP dose may be given as early as 12 months, if at least six months have passed since the 3rd DTaP dose was given.



Dear Colleague,

The Washington State Department of Health (DOH) provides print-ready files (PDFs) of health education materials. To ensure that the original quality of the piece is maintained, please read and follow the instructions below and the specifications included for professional printing.

- Use the latest version. DOH materials are developed using the most current information available, are checked for clinical accuracy, and are field tested with the intended audience to ensure they are clear and readable. DOH programs make periodic revisions to educational materials, so please check this web site to be sure you have the latest version. DOH assumes no responsibility for the use of this material or for any errors or omissions.
- **Do not alter**. We are providing this artwork with the understanding that it will be printed without alterations and copies will be free to the public. Do not edit the text or use illustrations or photographs for other purposes without first contacting us. Please do not alter or remove the DOH logo, publication number or revision date. If you want to use a part of this publication for other purposes, contact the Office of Health Promotion first.
- For quality reproduction: Low resolution PDF files are intended for black and white or color desktop printers. They work best if you are making only one or two copies. High resolution PDF files are intended for reproducing large quantities and are set up for use by professional offset print shops. The high resolution files also include detailed printing specifications. Please match them as closely as possible and insist on the best possible quality for all reproductions.

If you have questions, contact:
Office of Health Promotion
P.O. Box 47833 Olympia, WA 98504-7833
(360) 236-3736

Sincerely, Health Education Resource Exchange Web Team